



Nectar and Pollen Plants of Oklahoma

Phil Mulder

Area Extension Entomologist

There are few places in Oklahoma where surplus honey cannot be produced if bees are properly managed. However, locations only a few miles apart produce honey crops varying considerably in size. Therefore, greater profit might result if beekeepers give more attention to pollen and nectar sources near their apiaries. Some plants furnish only pollen and others only nectar, but the majority of plants supply both. Pollen is a source of protein, the tissue-building food for bees; nectar, a carbohydrate, is the energy food.

Experienced beekeepers have learned that in some years the ability of plants to produce nectar and pollen varies considerably. These variations may be the result of climatic conditions (droughts), changes in agricultural crops and prac-

Major Nectar and Pollen Plant Sources of Oklahoma

Plant Name*	Dates of Bloom
Alfalfa (N)	May 10 to Frost
Cotton (N&P)	July 10 to Frost
Hairy vetch (N&P)	April 20 to July 31
Persimmon (N&P)	May 15 to June 15
Plains sage (N&P)	June 1 to October
Smartweed (N&P)	July 1 to October
Spanishneedle (N&P)	Aug. 1 to Frost
Vitex (N&P)	May 15 to Frost
White sweetclover (N&P)	May 5 to July 25
Yellow sweetclover (N&P)	April 20 to July 10

Minor Nectar and Pollen Plant Sources of Oklahoma

Bee bee tree (N&P)	July 1 to August 31
Bindweed (N)	May 15 to Sept. 30
Black locust (N&P)	April 10 to May 20
Broomweed (N&P)	July 5 to Nov. 10
Dandelion (N&P)	April 1 to June 1
Elm (P)	Feb. 25 to March 25
Lawn clover (N&P)	March 20 to July 1
Lemon monarda (N&P)	June 1 to October
Maple (N&P)	Feb. 20 to May 5
Milkweed (N)	May 10 to Sept. 5
Pear (N&P)	March 15 to April 8
Redbud (N&P)	March 10 to April 15
Sand plum (N&P)	March 20 to May 10
Sumac (N&P)	May 10 to August 1
Sunflower (N&P)	July 1 to Frost
Tame plum (N&P)	March 20 to May 5
Willow (N&P)	March 20 to April 10

* N indicates nectar; P indicates pollen

Oklahoma Cooperative Extension Fact Sheets are also available on our website at: <http://osufacts.okstate.edu>

tices, damage by herbivorous insects, and the installation of irrigation projects.

Beekeepers should note when the relative abundance of plants important to the honey bee bloom, how long they remain in bloom, and the effect of weather conditions upon secretion of nectar.

Apiarists and farmers need information to aid them in coordinating honey production in Oklahoma with erosion control, pasture building, cover cropping, and the production of grain, fruit, vegetables, seeds, fiber, and livestock. The following is a list of important nectar and pollen plants of the state, and their dates of blooming.


Characteristics of Nectar and Pollen Plants

Major Sources


Alfalfa (*Medicago sativa*) is grown extensively over the entire state. Approximately one-third of honey produced in Oklahoma is from this source. Alfalfa honey is a very light amber to white with a reddish tint. A considerable amount of potential for honey production is not realized because alfalfa is primarily grown for hay and, during early summer, is harvested before full bloom. After July 1, however, farmers often stop cutting hay and leave alfalfa for seed production. The latter is compatible with honey production.

Cotton (*Gossypium hirsutum*) is one of the major crops of Oklahoma and is generally found in the southern one-half of the state. Cotton is an important nectar and pollen source when grown on rich alluvial soils. On lighter sandy soils, it produces little nectar. Cotton honey is white, has good body, and a characteristic cotton flavor. Because of the scarcity of bees in some cotton-growing areas and the frequent use of chemical insecticides, much potential for honey production is lost.

Hairy Vetch (*Vicia villosa*) produces an abundance of pollen and nectar. This plant may produce nectar for a surplus of honey in some areas of the state. One of the earliest major honey flows is obtained from this plant. The honey is water-white in color with good flavor and is well accepted by the consumer. Due to the early blooming period, bee colonies need to be very strong after the winter season to obtain a surplus from this plant.



Alfalfa



Sweetclover


Persimmon (*Diospyros virginiana*) produces large surpluses in the eastern one-third of the state. In fact, persimmon may be one of the state's major honey-producing plants. Honey produced from it is water-white.

Plains Sage (*Salvia coccinea*) produces a good source of nectar and a surplus of pollen. The honey is light in color, of good flavor, and has the characteristic monarda flavor. This plant is one of the major honey plants of Oklahoma and is found in abundance south of the Canadian River. Sometimes, falsely referred to as horsemint, this plant begins blooming the latter part of May or first of June.


Smartweed (*Polygonum* sp.) includes several species from the same genus. This plant grows statewide, usually as a volunteer weed, and produces honey that is dark in color with a characteristic buckwheat flavor and color. Most of this honey is used for wintering bees, but any surplus is readily used in bakeries or by the few people who appreciate the mild buckwheat taste.

Spanishneedle (*Bidens bipinnata* L.) is a dependable honey plant in northeastern Oklahoma providing a surplus of nectar. The honey is golden in color, with a good flavor.

Vitex (*Vitex negunda*) is also called summer lilac or chaste tree. This fast growing, multiple-stemmed shrub can reach a height of 12-15 feet. Aromatic whitish-purple spikes bloom all summer. This plant grows well in a wide variety of soil types. This plant is probably a major nectar and pollen source for apiaries in southern Oklahoma.



Hairy Vetch



smartweed


White sweetclover (*Melilotus alba*) grows well over the entire state and is also one of the major nectar and pollen producing plants of Oklahoma. It produces honey of good flavor, light color, and good body. White sweetclover blooms about 10-15 days after yellow sweetclover and produces a good surplus of pollen and nectar.

Yellow sweetclover (*Melilotus officinalis*) grows statewide and is a major source for honey production in Oklahoma. Nectar and pollen sources from yellow sweetclover tend to be 10-15 days ahead of white sweetclover, providing an earlier foraging site for honey bees. A surplus of honey may be obtained from this source; however, many colonies are not ready for the early nectar flow from this plant. Honey from yellow sweetclover is extra white to extra light amber with a mild flavor.

Minor Sources

Among the plants of lesser importance as sources of nectar and pollen in spring brood rearing are dandelion, elm, redbud, plum, willow, maple, locust, and fruit bloom.

Milkweed produces an abundance of pollen and, in some areas, produces a small surplus of nectar. This plant is found in the northeastern and north central counties of the state. Bindweed or tie vine is a very valuable honey plant in many sections of Oklahoma, especially on rich alluvial soils. This plant produces a surplus of white honey.



Spanish-needle



Persimmon

Sumac is found growing on thousands of acres in all sections of Oklahoma. It produces a surplus of light-colored and well-flavored honey. This plant and bindweed may become of less importance as herbicides are used to remove them from crops and pastures.

Broomweed is found growing statewide on prairies and is a good source of pollen and honey for wintering bees. A strong-flavored, golden-colored honey is usually gathered in September and October. Sunflower is also a good plant for producing pollen and nectar for fall brood rearing.

Oklahoma State University, in compliance with Title VI and VII of the Civil Rights Act of 1964, Executive Order 11246 as amended, Title IX of the Education Amendments of 1972, Americans with Disabilities Act of 1990, and other federal laws and regulations, does not discriminate on the basis of race, color, national origin, gender, age, religion, disability, or status as a veteran in any of its policies, practices, or procedures. This includes but is not limited to admissions, employment, financial aid, and educational services.

Issued in furtherance of Cooperative Extension work, acts of May 8 and June 30, 1914, in cooperation with the U.S. Department of Agriculture, Robert E. Whitson, Director of Cooperative Extension Service, Oklahoma State University, Stillwater, Oklahoma. This publication is printed and issued by Oklahoma State University as authorized by the Vice President, Dean, and Director of the Division of Agricultural Sciences and Natural Resources and has been prepared and distributed at a cost of 20 cents per copy. 0607